

PREPARATION FOR PROSTATE BIOPSY

If you are allergic to antibiotics or penicillin, contact us at 932 522 522 before starting the preparation.

MEDICATION:

Enema Casen 250cc (2)
Cefuroxima 500 mg (box of 14 pills)

BEFORE THE EXPLORATION:

- **If you take an antiaggregant therapy:** Aspirina, Adiro 300, AAS 300, Bioplak, Plavix, Clopidogrel... STOP 7 days BEFORE de la biopsy.
- **If you take an anticoagulant therapy:** Sintrom, acenocumarol, Aldocumar, Warfarina, Pradaxa, Dabigatran, Eliquis, Apixaban, Xarelto, Rivaroxaban GO TO YOUR DOCTOR to receive the medication suspension/restart guideline.
- The NIGHT BEFORE put ENEMA CASEN.
- The MORNING of the biopsy NOT EAT ANYTHING and take 1 pill of Cefuroxima 500 mg (with a little of water) and put the ENEMA CASEN.
- Come to the center with companion.
- It's forbidden drive until 24 hours before de exploration.

AFTER THE BIOPSY:

- Take 1 pill of Cefuroxima 500 mg each 12 hours until finish the box.
- You may bleed with urine, stool and semen for several days.
- If you have a fever over 38º go to emergencies.
- If you take antiaggregant, you can restart it the day after the test.
- If you take an anticoagulant, restart it according to the hemostasis guideline.